

US009428766B2

# (12) United States Patent

Goepfert et al.

## (10) Patent No.: US 9,428,766 B2

(45) **Date of Patent:** \*Aug. 30, 2016

# (54) PROTEIN EXPRESSION FROM MULTIPLE NUCLEIC ACIDS

## (71) Applicant: Hoffmann-La Roche Inc., Nutley, NJ

(US)

#### (72) Inventors: Ulrich Goepfert, Munich (DE);

Hendrik Knoetgen, Penzberg (DE); Erhard Kopetzki, Penzberg (DE); Anne Stern, Penzberg (DE)

#### (73) Assignee: HOFFMANN-LA ROCHE INC.,

Nutley, NJ (US)

## (\*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-

claimer.

#### (21) Appl. No.: 14/293,447

(22) Filed: Jun. 2, 2014

#### (65) Prior Publication Data

US 2014/0335609 A1 Nov. 13, 2014

#### Related U.S. Application Data

(63) Continuation of application No. 12/681,781, filed as application No. PCT/EP2008/008523 on Oct. 9, 2008, now Pat. No. 8,771,988.

#### (30) Foreign Application Priority Data

Oct. 12, 2007 (EP) ...... 07019999

(51)	Int. Cl.	
	C12P 21/08	(2006.01)
	C12N 15/85	(2006.01)
	C07K 16/18	(2006.01)
	C07K 16/28	(2006.01)
	C12P 21/00	(2006.01)

(52) **U.S. Cl.** 

### (58) Field of Classification Search

None

See application file for complete search history.

#### (56) References Cited

#### U.S. PATENT DOCUMENTS

5,420,019	A	5/1995	Theofan et al.
5,550,036	A	8/1996	Grinnell
5,639,275	A	6/1997	Baetge et al.
5,852,175	A	12/1998	Cummings et al.
2003/0096341	A1	5/2003	Mueller et al.
2005/0169925	A1	8/2005	Bardroff et al.
2005/0226876	A1	10/2005	Graus et al.

#### FOREIGN PATENT DOCUMENTS

0319206	6/1989
0921194	6/1996
1010758	6/2000
8900605	1/1989
89/10959 A	1 11/1989
93/01296 A	1 1/1993
94/25067 A	1 11/1994
95/17513 A	1 6/1995
99/47647 A	1 9/1999
0028066	5/2000
03/046187	6/2003
03070760	8/2003
2004/083247 A	1 9/2004
2004/087758 A	2 10/2004
	0921194 1010758 8900605 89/10959 A 93/01296 A 94/25067 A 95/17513 A 99/47647 A 0028066 03/046187 03070760 2004/083247 A

(Continued)

#### OTHER PUBLICATIONS

Jung, H. et al., Detection and treatment of mycoplasma contaminiation in cultured cells, Apr. 2003, CHang Gung Med. J., vol. 26, pp. 250-258.\*

The English translation of the Chinese Office Action, issued on Mar. 7, 2013, in the corresponding Chinese application No. 200880110216.6.

Summons to Oral Proceeding by the EPO, mailed on Oct. 8, 2014, in related European Patent No. 1453966.

(Continued)

Primary Examiner — Michael Burkhart (74) Attorney, Agent, or Firm — Jones Day

### (57) ABSTRACT

The current invention reports a method for the recombinant production of a secreted heterologous immunoglobulin in a CHO cell comprising the following steps: i) providing a CHO cell, which is adapted to growth in suspension culture, adapted to growth in serum-free medium, mycoplasma free. and virus free, ii) providing a vector comprising a prokaryotic origin of replication, a first nucleic acid conferring resistance to a prokaryotic selection agent, a second nucleic acid encoding the heavy chain of said heterologous immunoglobulin, a third nucleic acid encoding the light chain of said heterologous immunoglobulin, a fourth nucleic acid conferring resistance to a eukaryotic selection agent, iii) transfecting said CHO cell, wherein said transfecting comprises a) transfecting said CHO cell with said vector comprising a fourth nucleic acid conferring resistance to a first eukaryotic selection agent, b) selecting a CHO cell by growth in cultivation medium containing said first eukaryotic selection agent, c) transfecting said selected CHO cell with said vector comprising a fourth nucleic acid conferring resistance to a second eukaryotic selection agent different to said first eukaryotic selection agent, d) selecting a CHO cell by selected growth in cultivation medium containing said first and said second eukaryotic selection agent, iv) cultivating said transfected CHO cell in a medium in the presence of said first and second eukaryotic selection agent, under conditions suitable for the expression of said second, and third nucleic acid, and v) recovering said secreted heterologous immunoglobulin from the cultivation medium.